

<b>SF</b>	Mantle Inputs to the Subduction Factory: Assessing Scales of Spatial Variability along and Across the IBM Convergent Margin	
	Robert Stern, Univ. of Texas, Dallas; Bloomer; Clift	
	9/1/2001 – 8/31/2003	OCE 00-01827

Subduction Zones are complex physical and chemical systems where surficial Earth materials are dragged down to interact with asthenospheric mantle and generate melts. This interaction equilibrates Earth's mantle with its hydrosphere and atmosphere and produces continental crust, ore deposits, and explosive volcanism. Understanding how this interaction leads to melting, how these melts rise through downwelling mantle, and how melts evolve in the crust is a great scientific challenge, requiring multiple perspectives. One important perspective comes from understanding how the composition of magmas changes with depth to the Wadati-Benioff Zone (WBZ). From the earliest days of the plate tectonic revolution, systematic variations in magma chemistry—specifically, higher K contents with greater depth to the WBZ—were noted (Hatherton and Dickinson, 1969). In spite of this systematic behavior, it is still not clear what process is responsible. Hatherton and Dickinson (1969) were undecided as to its cause, suggesting that it could reflect either variations in the stability of K-bearing phases in the subducted slab (i.e., amphibole vs. phlogopite) or manifested differing ascent paths or melting processes.

This uncertainty persists. The 'K-h' relationship—since expanded to include more of the periodic table, including large ion lithophile (LIL) elements Rb and Ba and the light Rare Earths (LREE)—is still a robust constraint, but it is still not clear what this relationship reveals about subduction zone processes. We now know that the vast majority of major elements in arc melts come from partial melting of convecting asthenosphere in the mantle wedge, although incompatible element budgets may reflect distillation of subducted materials, especially sediments (Plank and Langmuir, 1993). Two attractive endmember possibilities are that the K-h relationship indicates that sediment melting becomes increasingly important with greater depth in the WBZ, or that lower degrees of melting of mantle peridotite occur as distance increases from the magmatic front.

Variations in sources and processes across the Subduction Factory can be most confidently addressed using volcanic cross-chains of intra-oceanic convergent margins (IOCM), those convergent margin systems that grow entirely within the ocean basins. IOCM magmatic systems may witness assimilation but it will be of relatively juvenile mantle lithosphere and thin mafic crust that are chemically and isotopically similar to the arc magmas that pass through, rather than the thicker, older, and more reactive crust and lithosphere associated with magmatic arcs on continents. IOCM cross-chains that extend rearward on to back-arc basin crust (like the Guguan Cross-Chain considered here) have minimal assimilation because these are built on crust that thins from ~20-30km along the magmatic front to ~5-10 km behind the arc. The Cook 7 expedition (March-April 2001) was funded by NSF-MARGINS for the purpose of addressing this issue. One of the best places to examine these variations is in a cross-chain associated with Guguan volcano in the Mariana arc system (Fig. 1).

The Guguan cross-chain extends west from the magmatic front, represented by Guguan island and a small submarine cone to the north (Cook 7 D50), which lie about 125 km above the WBZ. The West Guguan edifice (sampled by Cook 7 D49) lies on the western flank of Guguan, about 150 km above the WBZ. The Guguan 2 edifice lies over a part of the WBZ that is very steep, where we estimate a depth of 230km. Guguan is widely

acknowledged to be the endmember example of magma generation at convergent margins where the 'Subduction Component' is carried in an aqueous fluid (Elliott *et al.*, 1997). This interpretation agrees with observed high Ba/La and applies to N. Guguan as well (Fig. 2). Decreasing Ba/La and increasing La/Yb in lavas from W. Guguan and Guguan 2 could be interpreted as being due to the Subduction Component being carried as sediment melt, or could reflect low degrees of melting of the mantle. An explanation of sediment melting interpretation for the Guguan cross-chain is difficult to reconcile with Th/Nb and isotopic systematics. The isotopic variations reveal a mixing trend, but one which cannot simply be explained as a mixture between asthenospheric mantle and sediment melt. As is the case for the Izu cross-chains (Hochstaedter *et al.*, 2000), the rear-arc volcanoes in the Guguan cross-chain have a range of Th/Nb that is similar to that of the magmatic front, but at a lower  $^{87}\text{Sr}/^{86}\text{Sr}$ . Also like the Izu cross-chains, the Guguan cross-chain trends towards lower  $\epsilon_{\text{Nd}}$  and lower  $^{87}\text{Sr}/^{86}\text{Sr}$  with increasing depth to the subducted slab. The Guguan cross-chain also shows slightly lower  $\epsilon_{\text{Hf}}$  with increasing depth to the subducted slab. None of the trajectories seen in the isotopic variations are consistent with an interpretation that the rear-arc volcanoes were derived from melting of an asthenospheric mantle that had been 'pickled' in sediment melts. Two interpretations are consistent with the trace element and isotopic data: 1) Rear-arc melts result from mixtures of a "Guguan-like" metasomatized mantle with a hybrid MORB-type mantle that has been metasomatized by 1-2% sediment melt; or 2) Rear-arc melts result from melting of an enriched asthenospheric mantle, but with a role for fluids sufficient to cause arc-like enrichments in LIL elements. Work is continuing to address these possibilities, including trace element studies of melt inclusions, and Li and O isotopic compositions.

This information will also provide valuable constraints on thermal structure for OBS studies of upper mantle structure now in progress (triangles in Fig. 1B).

## Figures and Captions






Figure 1

**Figure 1:** A) Locality map of the Guguan Cross Chain in the Mariana arc system (courtesy Bob Embley, NOAA). B) The Central Mariana Arc system and the location of the Guguan Cross-Chain. Red triangles show the location of the U.S.-Japan passive land-OBS deployment carried out Summer 2003. C) HMR-1 sonar backscatter image of the Guguan cross-chain, showing the location of dredge sampling carried out during Cook 7 expedition. Note location of Guguan island. D) Cross-sectional view of the central Mariana subduction system using the earthquake catalog of Engdahl *et al.* (1988). Black circles denote earthquake hypocenters in a 50 km wide volume centered along 17.33°N.



**Figure 2:** Ba/La vs. La/Yb for lavas from the Guguan Cross-chain. Note that lavas erupted along the magmatic front (Guguan, N. Guguan) have higher Ba/La and lower La/Yb than those from W. Guguan and Guguan 2. The high Ba/La, low La/Yb component is MORB-type mantle that has been metasomatized by hydrous fluids. The low Ba/La, high La/Yb endmember could be sediment melts or lower degrees of melting of relatively enriched mantle. The composition of bulk sediment is shown, as is the composition of relatively Mariana Trough basalt associated with seafloor spreading.



**Figure 3:** Isotopic systematics of Guguan cross-chain lavas. A) Plot of Sr isotopes vs. Th/Nb. Curve shows fraction of sediment mixed with MORB-type asthenospheric mantle. High Th/Nb is taken to indicate melting of subducted sediments (Elliott *et al.*, 1997). Note that the range of Th/Nb encompassed by volcanoes along the magmatic front (Guguan and N. Guguan) is similar to that of the rear-arc volcanoes (W. Guguan and Guguan 2). These define parallel arrays but do not align along trajectories consistent with participation of sediment melts. B) Plot of Sr-Nd isotopes, with mixing trajectories for asthenospheric (MORB-type) mantle with bulk sediment compositions from ODP sites 800 and 801, east of the Mariana Trench (Plank and Langmuir, 1998). Note that the Guguan cross-chain samples are oriented perpendicular to the mixing trends. C) Plot of Sr-Hf isotopes (data for Guguan from Woodhead *et al.*, 2001), with mixing trajectories for asthenospheric (MORB-type) mantle with bulk sediment compositions from sediments east of the Mariana Trench (Plank and Langmuir, 1998; Woodhead, pers. comm. 2003). Note that the Guguan cross-chain samples are oriented perpendicular to the mixing trends.

## References

Elliott, T., Plank, T., Zindler, A., White, W. and Bourdon, B., 1997. Element transport from slab to volcanic front at the Mariana arc. *Journal of Geophysical Research*, B102: 14,991-15,019.

Engdahl, E.R., Hilst, R.D.v.d. and Buland, R., 1988. Global teleseismic earthquake relocation with improved travel times and procedures for depth determination. *Bulletin of the Seismological Society of America*, 88: 722-743.

Hatherton, T. and Dickinson, W.R., 1969. The relationship between andesitic volcanism and seismicity in Indonesia. *Journal of Geophysical Research*, 74(22): 5301-5310.

Hochstaedter, A.G., Gill, J.B., Taylor, B. and Ishizuka, O., 2000. Across-arc geochemical trends in the Izu-Bonin arc: Constraints on source composition and mantle melting. *Journal of Geophysical Research*, 105(B1): 495-512.

Ishihara, T., Stern, R.J., Fryer, P., Bloomer, S. and Becker, N.C., 2001. Seafloor spreading in the Southern Mariana Trough Inferred from 3-component Magnetometer Data. *EOS*, 82(47): F1202.

Kato, T. *et al.*, 2003. Geodetic evidence of back-arc spreading in the Mariana Trough. *Geophysical Research Letters*, 30(12): doi:10.1029/2002GL016757, 2003.

Plank, T. and Langmuir, C.H., 1993. Tracing trace elements from sediment input to volcanic output at subduction zones. *Nature*, 362: 739-742.

Plank, T., and Langmuir, C. H., 1998. The chemical composition of subducting sediment and its consequences for the crust and mantle, *Chem. Geol.*, 145, 325–394.

Woodhead, J.D., Hergt, J.M., Davidson, J.P., and Eggins, S.M., 2001. Hafnium isotope evidence for 'conservative' element mobility during subduction zone processes. *Earth and Planetary Science Letters* 192, 331-346.

## **Publications and Presentations**

### **Publications:**

Sun, C.-H., and Stern, R.J. 2001. Genesis of Mariana Shoshonites: Contribution of the 'Subduction Component' *Journal of Geophysical Research, Solid Earth*, v.106, 589-608.

Ohara, Y., Stern, R.J., Ishii, T., Yurimoto, H., and Yamazaki, T. 2002. Peridotites from the Mariana Trough: First look at the Mantle beneath an active Backarc Basin. *Contributions to Mineralogy and Petrology* v.143, 1-18.

Stern, R.J., 2002. "Subduction Zones" *Reviews of Geophysics*, 10.1029/2001RG000108

Ito, E., Stern, R.J., and Douthitt, C., 2003. 'Insights into Operation of the "Subduction Factory" from the Oxygen Isotopic Values of Southern Izu-Bonin-Mariana Arc' *The Island Arc* v. 12, 383-397.

Stern, R.J., Fouch, M.J., and Klempner, S., 2003. "An Overview of the Izu-Bonin-Mariana Subduction Factory" in J. Eiler and M. Hirschmann (eds.) *Inside the Subduction Factory*, *Geophysical Monograph* 138, American Geophysical Union, 175-222.

Gvirtzman, Z., and Stern, R.J., 2004. Bathymetry of Mariana Trench-Arc System and Formation of the Challenger Deep as a Consequence of Weak Plate Coupling. *Tectonics*, TC2011, doi:10.1029/2003TC001581, 2004

Stern, R.J., in press. Subduction Initiation: Induced and Spontaneous. *EPSL*

Kimura, J.-I., Stern, R.J., and Yoshida, T. in press. Re-initiation of Subduction and Magmatic Responses in SW Japan during Neogene Time. *Bull. Geol. Soc. America*

Kohut, E.J., Stern, R.J., Kent, A.J.R., Nielsen, R.L., Bloomer, S.H., and Leybourne, M. submitted. The Origins of High-Mg Lavas in the Mariana Arc: Evidence from Olivine-Hosted Melt Inclusions. *Journal of Petrology*.

**Presentations:**

Stern, R.J. "The Izu-Bonin-Mariana Subduction Factory" Keynote address and extended abstract for the Subduction Factory Theoretical and Experimental Institute, Eugene, OR (August 2000)

Kawate, S., Arima, M., Sun, C.H., Stern, R.J. 2000. "Crustal evolution in the Izu-Bonin-Mariana arc: New constraints from Pb-, Nd-, and Sr-isotopic data from the IBM Collision Zone, Central Honshu, Japan" EOS, v. 81, No. 48, F1247

Hargrove, U.S., Stern, R.J., Bloomer, S.H., Leybourne, M.I., Edwards, M., Becker, N. 2001. "Results of New High-Resolution Bathymetric and Back-scatter SONAR Imaging and Dredging of the Mariana Arc and Back-arc Basin System" GSA Abstract with Program v. 33, No. 6, A398

Stern, R.J., 2001. "Spontaneous Nucleation of Subduction Zones in the Western Pacific During Middle Eocene Time: Evidence from the IBM Forearc Ophiolite". EOS v. 84(47), F1225-1226.

Becker, N.C., Fryer, P., Martinez, F., Stern, R.J., Bloomer, S.H., 2001. "Magma Piracy in the Southern Mariana Backarc" EOS v. 84 (47), F1253

Bloomer, S.H., Stern, R.J., COOK 07 Shipboard Party 2001. "Mantle Inputs to the Subduction Factory: Detailed Studies of the Southern Mariana Seamount Province" EOS v. 84 (47), F1201-1202

Ishihara, T., Stern, R.J., Fryer, P., Bloomer, S.H., Becker, N.C., 2001. "Seafloor Spreading in the Southern Mariana Trough Inferred from 3-Component Magnetometer Data" EOS v. 84 (47), F1202

Bloomer, S.H., and Stern, R.J. 2002 "Tectonic Reorganization of the Western Pacific in Eocene Time: Missing Pieces in the Subduction Initiation Puzzle" EOS 83(47), F1251.

Stern, R.J., Hargrove, U.S., Leybourne, M. I. Pearce, J. A., and Bloomer, S.H., 2002 "An Overview of the Southern Mariana Subduction Factory: Arc, Cross-chains, and Back-arc Basin" EOS 83(47), F1317.

Stern, R.J., Bloomer, S.H., Leybourne, M., Hargrove III, U.S., Griffith, W.R., Fouch, M., Kohut, E, Vervoort, J., Kent, A. 2003. "Elemental Transport from the Descending Slab: Inferences from the Guguan Cross-Chain, Mariana Arc" State of the Arc Conference, Oregon

Kohut, Edward J., Stern, Robert J., Kent, Adam J., Nielsen, Roger L., and Bloomer, Sherman H. 2003. Discovery of Primitive 'Komatiitoid' lavas from the Mariana Arc Magmatic Front GSA Abstract with Program v. 35, No. 7, p. 182-183

Stern, R.J., and Hargrove, U.S. 2003. The Anatahan Felsic Province in the Mariana Arc System. Eos Trans. AGU 84(46), F1562.

Woodhead, J.D, Stern, R.J, Pearce, J. A. 2003. " Further Hf-Isotope Constraints on Element Mobility Within Subduction Zones" State of the Arc Conference, Oregon.

Wade, J.A., Plank, T., Stern, R.J., Hilton, D., Fischer, T., Moore, R., Trusdell, F., Saki, M., 2003. Geochemical Composition of Volcanic Rocks from the May 2003 Eruption of Anatahan Volcano, Mariana Islands. Eos Trans. AGU 84(46), F1562.

Kohut, E. J., Stern, R.J., Kent, A.J., Nielsen, R.L., and Bloomer, S.H. 2003. Across-Arc Variability of Outputs from the Mariana Subduction Factory: Melt Inclusions from the

14°35'N Cross-Chain, Eos Trans. AGU 84(46), F1398

Stern, R.J., Bloomer, S.H., Leybourne, M., Miller, N.R., Hargrove, U.S., Griffin, W.R., Fouch, M., Kohut, E., Vervoort, J., and Prytulak, J. 2003. Pickling peridotites in the IBM Mantle Wedge: Inferences from the Guguan Cross-chain, Mariana Arc. Eos Trans. AGU 84(46), F1398.

Shaw, A.M., Hauri, E.H., Hilton, D.R., Fischer, T.P., Stern, R.J., Wade, J., and Plank, T. 2004. Insights into arc fluid budgets from Mariana melt inclusions. 14<sup>th</sup> Goldschmidt Conference abstracts, p. A45.